

ON UNITS OF GROUP ALGEBRAS OF 2-GROUPS OF MAXIMAL CLASS

ZS. BALOGH, A. BOVDI

ABSTRACT. We determine the number of elements of order two in the group of normalized units $V(\mathbb{F}_2G)$ of the group algebra $\mathbb{F}_2G$ of a 2-group of maximal class over the field $\mathbb{F}_2$ of two elements. As a consequence for the 2-groups G and H of maximal class we have that $V(\mathbb{F}_2G)$ and $V(\mathbb{F}_2H)$ are isomorphic if and only if G and H are isomorphic.

Notation and results.

Let G be a finite p -group and $\mathbb{F}_pG$ its group algebra over the field $\mathbb{F}_p$ of p elements. For $x = \sum_{g \in G} \alpha_g g \in \mathbb{F}_pG$, $\alpha_g \in \mathbb{F}_p$ we denote $\sum_{g \in G} \alpha_g$ by $\chi(x)$. The subgroup

$$V(\mathbb{F}_pG) = \left\{ x = \sum_{g \in G} \alpha_g g \in \mathbb{F}_pG \mid \chi(x) = 1 \right\}$$

is called *the group of normalized units*. Evidently $V(\mathbb{F}_pG)$ is a p -group and its order is $p^{|G|-1}$.

Let $C = \langle a \mid a^{2^n} = 1 \rangle$ be the cyclic group of order 2^n , where $n \geq 2$. Consider the following extensions of C :

$$\begin{aligned} Q_{2^{n+1}} &= \langle a, b_1 \mid a^{2^n} = 1, b_1^2 = a^{2^{n-1}}, (a, b_1) = a^{-2}, n \geq 2 \rangle; \\ D_{2^{n+1}} &= \langle a, b_2 \mid a^{2^n} = 1, b_2^2 = 1, (a, b_2) = a^{-2}, n \geq 2 \rangle; \\ SD_{2^{n+1}} &= \langle a, b_3 \mid a^{2^n} = 1, b_3^2 = 1, (a, b_3) = a^{-2+2^{n-1}}, n \geq 3 \rangle, \end{aligned}$$

which are the generalized quaternion group, the dihedral group and the semidihedral group respectively. It is well-known that a finite 2-group of maximal class coincides with one of these groups.

2001 Mathematics Subject Classification. Primary 16A46, 16A26, 20C05. Secondary 19A22. Supported by Hungarian National Fund for Scientific Research (OTKA) grants No. T037202 and No. T043034

Our main result is the following:

Theorem. *Let G be a 2-group of maximal class and let $\Theta_G(2)$ be the number of elements of order two in $V(\mathbb{F}_2G)$. Then*

$$\begin{aligned}\Theta_{D_{2^{n+1}}}(2) &= 2^{2^n+n-1} + 2^{2^n}; \\ \Theta_{SD_{2^{n+1}}}(2) &= 2^{2^n+n-1}; \\ \Theta_{Q_{2^{n+1}}}(2) &= 2^{2^n+n-1} - 2^{2^n}.\end{aligned}$$

The following question is due to S. D. Berman: Let $\mathbb{F}_p$ be the field of p elements, G and H finite p -groups. Is it true that $V(\mathbb{F}_pG)$ and $V(\mathbb{F}_pH)$ are isomorphic if and only if G and H are isomorphic?

It is an interesting and likely more difficult question than the isomorphism problem for modular group algebras of p -groups over the field of p elements.

In [Berman] Berman gave a positive answer for this question for finite abelian p -groups. For nonabelian groups the question is open. As a consequence of the above theorem we have the

Corollary. *Let $\mathbb{F}_2$ be the field of 2 elements, and let G and H be finite 2-groups of maximal class. Then $V(\mathbb{F}_2G)$ is isomorphic to $V(\mathbb{F}_2H)$ if and only if G and H are isomorphic.*

Evidently if $V(\mathbb{F}_pG)$ and $V(\mathbb{F}_pH)$ are not isomorphic, then the group algebras $\mathbb{F}_pG$ and $\mathbb{F}_pH$ are not isomorphic. Moreover it follows from Baginski's result [1] that the group algebra $\mathbb{F}_2G$ of a 2-group G of maximal class uniquely determines the group G .

Some preliminary facts about group algebras of cyclic groups.

We observe some facts about abelian 2-groups and their group algebras.

Recall that for the group algebra $\mathbb{F}_2C$ of a cyclic 2-group $C = \langle a \mid a^{2^n} = 1 \rangle$ the subset

$$1, 1+a, (1+a)^2, \dots, (1+a)^{2^n-1} \tag{1}$$

is a basis of $\mathbb{F}_2C$ and a normalized unit u can be written as

$$1 + \sum_{i=1}^{2^n-1} \alpha_i (a+1)^i.$$

Each ideal of the group algebra $\mathbb{F}_2C$ has the form $\mathbb{F}_2C(1+a)^i$. Clearly $(1+a)^i$ belongs to the augmentation ideal $A(\mathbb{F}_2C)$ and by a Theorem of Jennings [6]

$$A^i(\mathbb{F}_2C) = A^{i+1}(\mathbb{F}_2C) + (a+1)^i \mathbb{F}_2.$$

According to [5] the annihilator of the element $(1+a)^i$ coincides with $A^{2^n-i}(\mathbb{F}_2C) = \mathbb{F}_2C(1+a)^{2^n-i}$.

Lemma 1. *The subgroup*

$$S_i = \{ \gamma \in V(\mathbb{F}_2 C) \mid \gamma(1+a)^i = (1+a)^i \}$$

of $V(\mathbb{F}_2 C)$ has order 2^i .

Proof. It is easy to see that $1 + \text{Ann}((1+a)^i)$ coincides with S_i , and according to [5] the cardinality $|\text{Ann}((1+a)^i)|$ is 2^i . □

Recall some well-known facts about a finite abelian 2-group N . Let $N[2]$ be the subgroup generated by elements of order two, and denote by $\Phi(N)$ the Frattini subgroup of N . We shall use the following

Lemma 2. *Let N be a finite abelian 2-group.*

1. *If $g \in \Phi(N)$, then $H_g = \{ h \in N \mid h^2 = g \}$ is a coset of N by $N[2]$.*
2. *If $g \notin \Phi(N)$, then $N = \langle g \rangle \times W$ for some subgroup W .*

Involutions and unitary subgroups.

Let $C = \langle a \mid a^{2^n} = 1, n \geq 2 \rangle$ be a cyclic 2-group. First we review some results on involutions of $\mathbb{F}_2 C$. Recall that the linear extension of the automorphism $a^i \rightarrow a^{-i}$ of G to the automorphism $x \rightarrow x^*$ of $\mathbb{F}_2 C$ is an involution of $\mathbb{F}_2 C$. Moreover, for $n \geq 3$ we have another involution $x \rightarrow x^\oplus$, which is generated by the automorphism $a^i \rightarrow a^{(2^{n-1}-1)i}$ of order two. For the sake of convenience we assume that σ is either $x \rightarrow x^*$ or $x \rightarrow x^\oplus$.

First we observe some properties of these involutions.

Lemma 3. *For $x = \alpha_0 + \alpha_1 a + \cdots + \alpha_{2^n-1} a^{2^n-1} \in \mathbb{F}_2 C$ we have*

$$x^2 = \sum_{i=0}^{2^{n-1}-1} (\alpha_i + \alpha_{i+2^{n-1}}) a^{2i}$$

and

$$xx^* = \chi(x) + \sum_{j=1}^{2^{n-1}-1} \left[\sum_{i=0}^{2^n-1} \alpha_i \alpha_{i-j \pmod{2^n}} \right] \cdot [a^j + a^{-j}].$$

Proof. Evidently $x^2 = \sum_{i=0}^{2^n-1} \alpha_i a^{2i}$ and $(a^{i+2^{n-1}})^2 = (a^i)^2$, thus the coefficient of a^{2i} equals $\alpha_i + \alpha_{i+2^{n-1}}$. Moreover, the equality $(xx^*)^* = xx^*$ asserts that

$$xx^* = \gamma_0 + \gamma_{2^{n-1}} a^{2^{n-1}} + \sum_{j=1}^{2^{n-1}-1} \gamma_j (a^j + a^{-j}).$$

For each $a^j \in \text{supp}(xx^*)$ with $0 \leq j \leq 2^{n-1} - 1$ the equality $a^j = a^i a^{-(i-j)}$ shows that $\gamma_j = \sum_{i=0}^{2^n-1} \alpha_i \alpha_{i-j} \pmod{2^n}$.

In particular, $\gamma_0 = \sum_{i=0}^{2^n-1} \alpha_i \alpha_i = \sum_{i=0}^{2^n-1} \alpha_i = \chi(x)$ and

$$\gamma_{2^{n-1}} = \sum_{i=0}^{2^n-1} \alpha_i \alpha_{i-2^{n-1}} \pmod{2^n} = 2 \cdot \sum_{i=0}^{2^{n-1}-1} \alpha_i \alpha_{i-2^{n-1}} \pmod{2^n} = 0. \quad \square$$

We put

$$Q = \{0, 2, 4, \dots, 2^{n-2} - 2\} \cup \{2^{n-1}, 2^{n-1} + 2, 2^{n-1} + 4, \dots, 2^{n-1} + 2^{n-2} - 2\},$$

$$P = \{0, 2, 4, \dots, 2^n - 2\} \text{ and } R = \{0, 2, 4, \dots, 2^{n-1} - 2\}.$$

Lemma 4. *For $x = \alpha_0 + \alpha_1 a + \dots + \alpha_{2^n-1} a^{2^n-1} \in \mathbb{F}_2 C$ we have*

$$xx^{\otimes} = \gamma_0 + \gamma_{2^{n-1}} a^{2^{n-1}} + \sum_{k \in R \setminus \{0\}} \gamma_k (a^k + a^{-k}) + \sum_{k \in 1+Q} \gamma_k (a^k + a^{-k+2^{n-1}}),$$

where

$$\gamma_k = \begin{cases} \sum_{r \in P} \alpha_r \alpha_{r-k} \pmod{2^n} + \sum_{r \in 1+P} \alpha_r \alpha_{r-k+2^{n-1}} \pmod{2^n} & \text{for } k \in P \setminus \{0, 2^{n-1}\}; \\ \sum_{r \in P} \alpha_r \alpha_{r-k+2^{n-1}} \pmod{2^n} + \sum_{r \in 1+P} \alpha_r \alpha_{r-k} \pmod{2^n} & \text{for } k \in 1+P; \\ \sum_{r \in 1+P} \alpha_r & \text{for } k = 2^{n-1}; \\ \sum_{r \in P} \alpha_r & \text{for } k = 0. \end{cases}$$

Proof. Since $(xx^{\otimes})$ is $\otimes$ -symmetric so

$$xx^{\otimes} = \gamma_0 + \gamma_{2^{n-1}} a^{2^{n-1}} + \sum_{k \in R \setminus \{0\}} \gamma_k (a^k + a^{-k}) + \sum_{k \in 1+Q} \gamma_k (a^k + a^{-k+2^{n-1}}).$$

We define the permutation ρ of the set $\{0, 1, 2, \dots, 2^n - 1\}$ in the following way:

$$\rho(i) = \begin{cases} i & \text{if } i \text{ is even;} \\ i + 2^{n-1} \pmod{2^n} & \text{if } i \text{ is odd.} \end{cases}$$

Using the permutation ρ , simple computations show that $x^{\otimes} = \sum_{i=0}^{2^n-1} \alpha_{\rho(i)} a^{-i}$ and the coefficient of $a^k \in \text{supp}(xx^{\otimes})$ is equal to the trace $\gamma_k = \text{tr}(xx^{\otimes} a^{-k})$. Evidently

$$\gamma_k = \text{tr}\left(\left(\sum_{j=0}^{2^n-1} \alpha_j a^j\right) \cdot \left(\sum_{i=0}^{2^n-1} \alpha_{\rho(i-k)} a^{-i}\right)\right) = \sum_{r=0}^{2^n-1} \alpha_r \alpha_{\rho(r-k)}.$$

Therefore we have for even k

$$\gamma_k = \sum_{r=0}^{2^n-1} \alpha_r \alpha_{\rho(r-k)} = \sum_{r \in P} \alpha_r \alpha_{r-k \pmod{2^n}} + \sum_{r \in 1+P} \alpha_r \alpha_{r-k+2^{n-1} \pmod{2^n}},$$

and for odd k

$$\gamma_k = \sum_{i \in P} \alpha_i \alpha_{i-k+2^{n-1} \pmod{2^n}} + \sum_{i \in 1+P} \alpha_i \alpha_{i-k \pmod{2^n}}.$$

Specifically for $k = 2^{n-1}$ simple computations show that

$$\gamma_{2^{n-1}} = \sum_{i \in P} \alpha_i \alpha_{i-2^{n-1} \pmod{2^n}} + \sum_{i \in 1+P} \alpha_i \alpha_i = \sum_{i \in 1+P} \alpha_i,$$

and

$$\gamma_0 = \sum_{i \in P} \alpha_i \alpha_i + \sum_{i \in 1+P} \alpha_i \alpha_{i-2^{n-1} \pmod{2^n}} = \sum_{i \in P} \alpha_i.$$

□

Recall that each involution σ of $\mathbb{F}_2 C$ determines a σ -unitary subgroup

$$V_\sigma(\mathbb{F}_2 C) = \{ \gamma \in V(\mathbb{F}_2 C) \mid \gamma^{-1} = \gamma^\sigma \}$$

of $V(\mathbb{F}_2 C)$.

The structure of $*$ -unitary subgroup of $V(\mathbb{F}_2 C)$ was described in [3] and [4]. According to these results the order of $V_*(\mathbb{F}_2 C)$ is

$$|C^2[2]| \cdot |\mathbb{F}_2|^{\frac{1}{2}(|C|+|C[2]|)-1} = 2^{2^{n-1}+1}. \quad (2)$$

Now we determine the order of the $\otimes$ -unitary subgroup $V_\otimes(\mathbb{F}_2 C)$, and as far as we know, it has not been investigated so far.

The mapping φ_σ , given by

$$\varphi_\sigma(x) = x^\sigma x^{-1} \quad \text{for } x \in V(\mathbb{F}_2 C),$$

is a homomorphism of the group $V(\mathbb{F}_2 C)$ onto some subgroup $W_\sigma(C)$ of the σ -unitary subgroup $V_\sigma(\mathbb{F}_2 C)$ with kernel $S_\sigma(C) = \{ x \in V(\mathbb{F}_2 C) \mid x = x^\sigma \}$.

The subset D of C determines the element $\hat{D} = \sum_{g \in D} g$ of $F_2 C$.

Lemma 5. *The unit $1 + \widehat{C}$ does not belong to $W_{\otimes}(C)$ and $1 + \widehat{C}^2$ is not an element of the subgroup $V_{\otimes}^2(\mathbb{F}_2C)$.*

Proof. Assume that $1 + \widehat{C} \in W_{\otimes}(C)$. Then $1 + \widehat{C} = x^{\otimes}x^{-1}$ for some $x \in V(\mathbb{F}_2C)$ and

$$x^{\otimes} = x(1 + \widehat{C}) = x + \chi(x)\widehat{C} = x + \widehat{C}. \quad (3)$$

Evidently the traces of the elements x and $x^{\otimes}$ are equal, so $\text{tr}(x + x^{\otimes}) = 0$ and the equality (3) leads to contradiction at characteristic two.

Now, suppose that $1 + \widehat{C}^2 = x^2$ for some $\otimes$ -unitary element $x \in V_{\otimes}(\mathbb{F}_2C)$. Since $x^{\otimes} = x^{-1}$ and x has the form $x = y_1 + y_2a$ with $y_i \in \mathbb{F}_2C^2$, so

$$x^{\otimes} + x = x(1 + x^{-2}) = x(1 + x^2) = \widehat{C}^2(\chi(y_1) + \chi(y_2)a^{\otimes}),$$

and $\chi(y_1) = \text{tr}(x^{\otimes} + x) = 0$. This shows that $y_1^{\otimes} + y_1 + y_2a + (y_2a)^{\otimes} = \widehat{C}^2a^{\otimes}$, which implies that $y_1 = y_1^{\otimes}$ and

$$\widehat{C}^2 = y_2^{\otimes} + y_2a^{2^{n-1}+2}.$$

The previous equality shows that the set $\text{supp}(y_2^{\otimes}) \cap \text{supp}(y_2a^{2^{n-1}+2})$ is empty, so $|\text{supp}(y_2^{\otimes})| = 2^{n-2}$, thus $\chi(y_2) = 0$, a contradiction. $\square$

The next homomorphism of $V(\mathbb{F}_2C)$ we shall use later. Evidently if $\Psi_{\sigma}(x) = xx^{\sigma}$ for $x \in V(\mathbb{F}_2C)$, then Ψ_{σ} is a homomorphism of $V(\mathbb{F}_2C)$ with kernel $V_{\sigma}(\mathbb{F}_2C)$.

Lemma 6. *If $C = \langle a \mid a^{2^n} = 1, n \geq 3 \rangle$, then the order of $V_{\otimes}(\mathbb{F}_2C)$ is $2^{\frac{|C|}{2}}$ and*

$$V_{\otimes}(\mathbb{F}_2C) = W_{\otimes}(C) \times \langle 1 + \widehat{C} \rangle.$$

Proof. It is easy to see that the lower layers of the both groups $V_{\otimes}(\mathbb{F}_2C)$ and $S_{\otimes}(C)$ coincide. To determine the 2-rank of $V_{\otimes}(\mathbb{F}_2C)$ it suffices to find the order of the lower layer $S_{\otimes}(C)[2]$ of the group $S_{\otimes}(C)$.

Each $\otimes$ -symmetric element of $V(\mathbb{F}_2C)$ has the form

$$\gamma_0 + \gamma_{2^{n-1}}a^{2^{n-1}} + \sum_{k \in R \setminus \{0\}} \gamma_k(a^k + a^{-k}) + \sum_{k \in 1+Q} \gamma_k(a^k + a^{-k+2^{n-1}}),$$

where $\gamma_i \in \mathbb{F}_2$ and $\gamma_0 + \gamma_{2^{n-1}} = 1$. Therefore $|S_{\otimes}(C)| = 2^{\frac{|C|}{2}}$ and analogously, $|S_{\otimes}(C)^2| = 2^{\frac{|C^2|}{2}-1}$. This asserts that the order of the subgroups $S_{\otimes}(C)[2]$ and $V_{\otimes}(\mathbb{F}_2C)[2]$ is $2^{\frac{|C|}{4}+1}$. But the kernel of $\varphi_{\otimes}$ is $S_{\otimes}(C)$, so

$$|W_{\otimes}(C)| = |V(\mathbb{F}_2C)| : |S_{\otimes}(C)| = 2^{\frac{|C|}{2}-1}.$$

Now we are ready to prove by induction on the order of C that $|V_{\oplus}(\mathbb{F}_2 C)| = 2^{\frac{|C|}{2}}$. First let $|C| = 2^3$. According to Lemma 4 for the unit $x = \sum_{i=0}^7 \alpha_i a^i$ we have

$$xx^{\oplus} = \beta_0 + \beta_1(a + a^3 + a^5 + a^7) + \beta_2(a^2 + a^6) + (\beta_0 + 1)a^4,$$

because $\chi(x) = 1$.

Since $\psi_{\oplus}(x) = xx^{\oplus}$, and the order of $Im(\psi_{\oplus})$ is 2^3 , so the order of $V_{\oplus}(\mathbb{F}_2 C)$ is

$$|V(\mathbb{F}_2 C)| : |Im(\psi_{\oplus})| = 2^4.$$

Evidently $1 + \widehat{C} \in V_{\oplus}(\mathbb{F}_2 C)$ and $1 + \widehat{C} \notin W_{\oplus}(C)$ by Lemma 5. Since the subgroup $\langle 1 + \widehat{C} \rangle \times W_{\oplus}(C)$ of $V_{\oplus}(\mathbb{F}_2 C)$ has order 2^4 thus

$$V_{\oplus}(\mathbb{F}_2 C) = \langle 1 + \widehat{C} \rangle \times W_{\oplus}(C).$$

Now let $|C| > 2^3$. Applying the inductive hypothesis for C^2 ,

$$V_{\oplus}(\mathbb{F}_2 C^2) = W(C^2) \times \langle 1 + \widehat{C^2} \rangle.$$

Lemma 5 asserts that $\langle 1 + \widehat{C} \rangle \times W_{\oplus}(C)$ is a subgroup of $V_{\oplus}(\mathbb{F}_2 C)$ and $\langle 1 + \widehat{C^2} \rangle \times V_{\oplus}^2(\mathbb{F}_2 C)$ is a subgroup of $V_{\oplus}(\mathbb{F}_2 C^2)$. This shows that

$$|V_{\oplus}^2(\mathbb{F}_2 C)| \leq |V_{\oplus}(\mathbb{F}_2 C^2)| : |\langle 1 + \widehat{C^2} \rangle| = |W(C^2)|,$$

so

$$|V_{\oplus}(\mathbb{F}_2 C)| = |V_{\oplus}^2(\mathbb{F}_2 C)| \cdot |V_{\oplus}(\mathbb{F}_2 C)[2]| \leq |W(C^2)| \cdot |V_{\oplus}(\mathbb{F}_2 C)[2]| = 2^{\frac{|C|}{2}}.$$

But $\langle 1 + \widehat{C} \rangle \times W_{\oplus}(C)$ is a subgroup of $V_{\oplus}(\mathbb{F}_2 C)$ and its order is $2^{\frac{|C|}{2}}$, therefore the order of $V_{\oplus}(\mathbb{F}_2 C) = \langle 1 + \widehat{C} \rangle \times W_{\oplus}(C)$ is $2^{\frac{|C|}{2}}$. □

Elements of order two in $V(\mathbb{F}_2 G)$.

For a fixed noninvertible element $z \in \mathbb{F}_2 C$ the set

$$M_z^{\sigma} = \{ y \in V(\mathbb{F}_2 C) \mid (y + y^{\sigma})z = 0 \}$$

is a subgroup of $V(\mathbb{F}_2 C)$. Indeed, if $y_1, y_2 \in M_z^{\sigma}$, then $y_1 z = y_1^{\sigma} z$ and $y_2 z = y_2^{\sigma} z$. Hence $y_1 y_2 z = y_1 y_2^{\sigma} z = y_1^{\sigma} y_2^{\sigma} z = (y_1 y_2)^{\sigma} z$, so $y_1 y_2 \in M_z^{\sigma}$.

In $V(\mathbb{F}_2 G)$ we divide the units of order two into two classes. It is well-known that $x = x_1 + x_2 b$ is a unit if and only if $\chi(x_1) + \chi(x_2) = 1$.

Definition. Let $x = x_1 + x_2b$ be a unit. If x has order two, $\chi(x_1) = 1$ and $\chi(x_2) = 0$, then x is called a unit of type 1. If x has order two, $\chi(x_1) = 0$ and $\chi(x_2) = 1$, then x is called a unit of type 2.

Lemma 7. The number of units of type 1 in both groups $V(F_2D_{2^{n+1}})$ and $V(F_2Q_{2^{n+1}})$ is equal.

Proof. A unit $x_1 + x_2b_2 \in V(\mathbb{F}_2D_{2^{n+1}})$ has order two if and only if

$$\begin{cases} x_1^2 = x_2x_2^* + 1; \\ (x_1 + x_1^*)x_2 = 0. \end{cases} \quad (4)$$

Similarly, $x_1 + x_2b_1 \in V(\mathbb{F}_2Q_{2^{n+1}})$ is a unit of order two if and only if

$$\begin{cases} x_1^2 = x_2x_2^*a^{2^{n-1}} + 1; \\ (x_1 + x_1^*)x_2 = 0. \end{cases}$$

Let $x_2 \in \mathbb{F}_2C$ be a fixed not invertible element. Clearly $x_2x_2^* + 1$, $x_2x_2^*a^{2^{n-1}} + 1$ belong to the subgroup M_{x_2} , defined before, and the set

$$H_{x_2x_2^*+1} = \{ h \in M_{x_2}^* \mid h^2 = x_2x_2^* + 1 \}$$

either is empty or according to Lemma 2 constitutes a coset of $M_{x_2}^*$ by $M_{x_2}^*[2]$. Similarly,

$$H_{x_2x_2^*a^{2^{n-1}}+1} = \{ h \in M_{x_2}^* \mid h^2 = x_2x_2^*a^{2^{n-1}} + 1 \}$$

is either a coset of $M_{x_2}^*$ by $M_{x_2}^*[2]$ or an empty set.

Let us prove that $x_2x_2^* + 1 \in \Phi(M_{x_2}^*)$ if and only if $x_2x_2^*a^{2^{n-1}} + 1 \in \Phi(M_{x_2}^*)$. Suppose that $u = x_2x_2^* + 1 \notin \Phi(M_{x_2}^*)$ and $v = x_2x_2^*a^{2^{n-1}} + 1 \in \Phi(M_{x_2}^*)$. Then for $u \notin \Phi(M_{x_2}^*)$ there exists a direct decomposition $M_{x_2}^* = \langle u \rangle \times W$ for some subgroup W . Of course

$$v^2 = (x_2x_2^*a^{2^{n-1}} + 1)^2 = (x_2x_2^*)^2 + 1 = u^2,$$

therefore $u^{-1}v \in M_{x_2}^*[2]$ and $v = uy$ for some $y \in M_{x_2}^*[2]$. Obviously, $v \in \Phi(M_{x_2}^*) = \langle u^2 \rangle \times W^2$, whence $v = u^{2t}w^2$ for some t and $w \in W$. Moreover, $v = uy$ implies that

$$y = u^{2t-1}w^2 \in M_{x_2}^*[2] \quad \text{and} \quad 1 = y^2 = u^{4t-2}w^4.$$

This shows that $u^{4t-2} = 1$, so $4t \equiv 2 \pmod{|u|}$, where $|u|$ is the order of u . Hence $|u| = 2$ and $(x_2x_2^*)^2 = 0$. By Lemma 3, $x_2x_2^* = z(1 + a^{2^{n-1}})$ for some $z \in \mathbb{F}_2C$ and

$$v = x_2x_2^*a^{2^{n-1}} + 1 = z(1 + a^{2^{n-1}})a^{2^{n-1}} + 1 = z(1 + a^{2^{n-1}}) + 1 = x_2x_2^* + 1 = u$$

which is impossible.

We have shown that the cardinalities of the subsets $H_{x_2x_2^*+1}$ and $H_{x_2x_2^*a^{2^{n-1}}+1}$ are equal for each not invertible element x_2 , thus the proof of lemma is complete. $\square$

A unit $x_1 + x_2b_3 \in V(\mathbb{F}_2SD_{2^{n+1}})$ has order two if and only if

$$\begin{cases} x_1^2 = x_2x_2^{\circledast} + 1; \\ (x_1 + x_1^{\circledast})x_2 = 0. \end{cases} \quad (5)$$

For each noninvertible and nonzero element x_2 there exists an i such that $x_2 = \gamma(1+a)^i$ for some $\gamma \in V(\mathbb{F}_2C)$. The equalities (4) and (5) assert that

$$x_2x_2^* = \gamma\gamma^*(1+a)^{2i}a^{-i} \in \mathbb{F}_2C^2,$$

and

$$x_2x_2^{\circledast} = \gamma\gamma^{\circledast}(1+a)^i(1+a^{2^{n-1}-1})^i \in \mathbb{F}_2C^2.$$

In particular, if $i = 2l$ is even, then

$$(1+a)^{2l}(1+a^{2^{n-1}-1})^{2l} = (1+a^2)^l(1+a^{-2})^l = (1+a)^{2l}(1+a^{-1})^{2l},$$

and $x_2x_2^{\circledast} = \gamma\gamma^{\circledast}(1+a)^{2i}a^{-i}$.

For each $0 \leq i < 2^n$ we define the set

$$H_i^\sigma = \{ h \in V(\mathbb{F}_2C) \mid hh^\sigma(1+a)^i(1+a^\sigma)^i \in \mathbb{F}_2C^2 \}.$$

Lemma 8. *The set H_i^σ has the following properties:*

1. *If $i \geq 2^{n-1}$, then $H_i^\sigma = V(\mathbb{F}_2C)$.*
2. *If $i = 2l + 1$ is odd and $i < 2^{n-1}$, then the set H_i^σ is empty.*
3. *If $i = 2l$ is even and $i < 2^{n-1}$, then H_{2l}^σ is a subgroup of $V(\mathbb{F}_2C)$.*
4. *For even indices $i = 2l < 2^{n-1}$*

$$H_0^\sigma \subset H_2^\sigma \subset H_4^\sigma \subset \cdots \subset H_{2l}^\sigma \subset \cdots \subset H_{2^{n-1}-2}^\sigma = V(\mathbb{F}_2C)$$

and the order of the subgroup H_{2l}^σ is $2^{3 \cdot 2^{n-2} + l}$.

Proof. 1. Since $(1+a)^{2^{n-1}} = (1+a^\sigma)^{2^{n-1}}$, for $i \geq 2^{n-1}$ we have

$$hh^\sigma(1+a)^i(1+a^\sigma)^i = hh^\sigma(1+a)^{2^n}(1+a)^{i-2^{n-1}}(1+a^\sigma)^{i-2^{n-1}} = 0.$$

Thus $hh^\sigma(1+a)^i(1+a^\sigma)^i \in \mathbb{F}_2C^2$ for each $h \in V(\mathbb{F}_2C)$. Hence $H_i^\sigma = V(\mathbb{F}_2C)$ for all $i \geq 2^{n-1}$.

2. Let $i = 2l + 1$ be a fixed odd integer and $i < 2^{n-1}$.

First, let us prove by induction on l that if $i = 2l + 1$, then

$$(a + a^{-1})^i = \sum_{r \text{ is odd}} \beta_r (a^r + a^{-r}), \quad (6)$$

where $\beta_r \in \mathbb{F}_2$. It is clear for $l = 0$ and assume that

$$(a + a^{-1})^{2l+1} = \sum_{r \text{ is odd}} \beta_r (a^r + a^{-r}).$$

By the identity

$$(a^s + a^{-s})(a^k + a^{-k}) = (a^{s+k} + a^{-(s+k)}) + (a^{s-k} + a^{-(s-k)}) \quad (7)$$

we have

$$(a + a^{-1})^{2(l+1)+1} = \sum_{r \text{ is odd}} \beta_r ((a^{r+2} + a^{-r-2}) + (a^{r-2} + a^{-r+2}))$$

the desired assertion.

Now, we begin to prove that the set H_i^σ is empty. First let σ be the $*$ -involution. By Lemma 3, for $h \in H_i^*$ we have $hh^* = 1 + z_1 + z_2$, where

$$z_1 = \sum_{j \in R \setminus \{0\}} \alpha_j (a^j + a^{-j}), \quad z_2 = \sum_{k \in 1+R} \alpha_k (a^k + a^{-k}),$$

and $R = \{0, 2, 4, \dots, 2^{n-1} - 2\}$. By definition,

$$hh^*(a + a^{-1})^i = (1 + z_1)(a + a^{-1})^i + z_2(a + a^{-1})^i \in \mathbb{F}_2 C^2 \quad (8)$$

and since $i = 2l + 1$ is odd it is easy to see that

$$(a^k + a^{-k})(a + a^{-1})^i = (a^k + a^{-k})(a + a^{-1})(a^2 + a^{-2})^l,$$

so by (6) and (7) we have $z_2(a + a^{-1})^i \in \mathbb{F}_2 C^2$ and $w = (1 + z_1)(a + a^{-1})^i \notin \mathbb{F}_2 C^2$ if $w \neq 0$. But (8) confirms that $(1 + z_1)(a + a^{-1})^i = 0$ and we have $(a + a^{-1})^i = 0$, because $1 + z_1$ is a unit. This is impossible for $i < 2^{n-1}$.

Let σ be the $\otimes$ -involution. Lemma 4 asserts that $hh^\otimes = 1 + z_1 + z_2$, where

$$z_1 = (\gamma_0 + 1)(1 + a^{2^{n-1}}) + \sum_{k \in R \setminus \{0\}} \gamma_k (a^k + a^{-k}),$$

$$z_2 = \sum_{k \in 1+Q} \gamma_k (a^k + a^{-k+2^{n-1}})$$

and

$$Q = \{ 0, 2, 4, \dots, 2^{n-2} - 2 \} \cup \{ 2^{n-1}, 2^{n-1} + 2, 2^{n-1} + 4, \dots, 2^{n-1} + 2^{n-2} - 2 \}.$$

We remark that

$$\begin{aligned} & hh^{\otimes}(1+a)^{2l+1}(1+a^{2^{n-1}-1})^{2l+1} = \\ & = (1+z_1+z_2)(a+a^{-1})^{2l} \left((1+a)^{2^{n-1}} + (a+a^{-1}) + (1+a)^{2^{n-1}} a^{-1} \right) \in \mathbb{F}_2 C^2. \end{aligned}$$

The identity

$$(a^k + a^{-k+2^{n-1}})(a^r + a^{-r}) = (a^{k+r} + a^{-r-k+2^{n-1}}) + (a^{k-r} + a^{-k+r+2^{n-1}})$$

and (7) assert that

$$(1+z_1)(a+a^{-1})^{2l}(1+a)^{2^{n-1}} + z_2(a+a^{-1})^{2l}(a+a^{-1}) + z_2(a+a^{-1})^{2l}(1+a)^{2^{n-1}} a^{-1}$$

belongs to $\mathbb{F}_2 C^2$, and

$$w = (1+z_1)(a+a^{-1})^{2l+1} + (a+a^{-1})^{2l}(1+a)^{2^{n-1}} z_2 + (a+a^{-1})^{2l}(1+a)^{2^{n-1}} (1+z_1)a^{-1}$$

is not an element of $\mathbb{F}_2 C^2$, if $w \neq 0$. By (8), $w \in \mathbb{F}_2 C^2$, but this leads again to a contradiction. Therefore

$$(1+z_1)(a+a^{-1})^{2l+1} + (a+a^{-1})^{2l}(1+a)^{2^{n-1}} (z_2 + (1+z_1)a^{-1}) = 0.$$

Since $1+z_1$ and $e = z_2(1+z_1)^{-1} + a^{-1}$ are units, it follows that

$$(a+a^{-1})^{2l} \left((a+a^{-1}) + (1+a)^{2^{n-1}} e \right) = 0. \quad (9)$$

For $l = 0$ we have $(a+a^{-1}) = (1+a)^{2^{n-1}} e$, which is impossible, because $(a+a^{-1})^2 \neq 0$. If $l > 0$, then

$$(a+a^{-1})^{2l}(1+a)^{2^{n-1}} e \in A^{4l+2^{n-1}}(\mathbb{F}_2 C).$$

Thus from (9) it follows that $(a+a^{-1})^i \in A^{4l+2^{n-1}}(\mathbb{F}_2 C)$ and $i < 4l + 2^{n-1}$. This is impossible by Jennings's theory [4]. Thus H_i^σ is empty, as we stated.

3. Now we shall prove that for $i = 2l < 2^{n-1}$ the set H_i^σ is a subgroup of $V(\mathbb{F}_2 C)$. For $h \in H_i^\sigma$ using the basis (1) we have

$$hh^\sigma = 1 + \sum_{j=1}^{2^{n-1}-1} \alpha_j (1+a)^j,$$

and

$$hh^\sigma(1+a)^{2i} = (1+a)^{2i} + \sum_{j=1}^{2^n-2i-1} \alpha_j(1+a)^{j+2i} \in \mathbb{F}_2C^2.$$

But it follows that $\alpha_1 = \alpha_3 = \alpha_5 = \cdots = \alpha_{2^n-2i-1} = 0$. Consequently, for $h_1, h_2 \in H_i^\sigma$ there exist $u_1, u_2 \in \mathbb{F}_2C^2$ and $v_1, v_2 \in A^{2^n-2i}(\mathbb{F}_2C)$ such that

$$h_1h_1^\sigma = 1 + u_1 + v_1 \quad \text{and} \quad h_2h_2^\sigma = 1 + u_2 + v_2.$$

Clearly

$$h_1h_2(h_1h_2)^\sigma = h_1h_1^\sigma h_2h_2^\sigma = 1 + u_1 + u_2 + u_1u_2 + z,$$

where $1 + u_1 + u_2 + u_1u_2 \in \mathbb{F}_2C^2$ and $z = v_1 + v_1u_2 + v_2 + u_1v_2 + v_1v_2 \in A^{2^n-2i}(\mathbb{F}_2C)$. Hence $h_1h_2 \in H_i^\sigma$ and H_i^σ is a subgroup of $V(\mathbb{F}_2C)$.

In the next step we shall verify that $H_{2^{n-1}-2}^\sigma = V(\mathbb{F}_2C)$. A unit u of $\mathbb{F}_2C$ can be written as

$$u \equiv 1 + \alpha_1(1+a) + \alpha_2(1+a)^2 + \alpha_3(1+a)^3 \pmod{A^4(\mathbb{F}_2C)}.$$

It is easy to see that

$$\begin{aligned} (1+a^\sigma) &\equiv (1+a) + (1+a)^2 + \\ &\quad + (1+a)^3 + \cdots + (1+a)^{2^{n-1}-1} \pmod{A^{2^{n-1}}(\mathbb{F}_2C)}. \end{aligned} \quad (10)$$

It follows that

$$u^\sigma \equiv 1 + \alpha_1(1+a) + (\alpha_1 + \alpha_2)(1+a)^2 + (\alpha_1 + \alpha_3)(1+a)^3 \pmod{A^4(\mathbb{F}_2C)}$$

and

$$uu^\sigma \equiv 1 \pmod{A^4(\mathbb{F}_2C)}. \quad (11)$$

Therefore for each $h \in V(\mathbb{F}_2C)$ and $i = 2^{n-1} - 2$, (11) confirms that

$$hh^\sigma(1+a)^{2i} = hh^\sigma(1+a)^{(2^n-4)} = (1+a)^{(2^n-4)}.$$

This shows that $h \in H_{2^{n-1}-2}^\sigma$ and $H_{2^{n-1}-2}^\sigma = V(\mathbb{F}_2C)$, because h is an arbitrary unit. This completes the proof of this assertion.

4. Now let $i = 2l < 2^{n-1} - 2$. We shall prove that H_i^σ is a proper subgroup of H_{i+2}^σ . Clearly $H_i^\sigma \subseteq H_{i+2}^\sigma$ and it is sufficient to verify that $h = 1 + (1+a)^{2^n-(2i+3)}$ does not belong to H_i^σ and $h \in H_{i+2}^\sigma$.

Note that for an even i the binomial coefficient $\binom{2^n-2i-3}{2}$ is even too and $h^\sigma = 1 + (1 + a^\sigma)^{2^n-(2i+3)}$. By the binomial formula and (10) we have

$$h^\sigma = 1 + (1 + a)^{2^n-(2i+3)} + (1 + a)^{2^n-(2i+2)} + (1 + a)^{2^n-(2i+1)} + x_\sigma,$$

where $x_\sigma \in A^{2^n-2i}(\mathbb{F}_2C)$. It is easy to see that $2^{n+1}-4i-j \geq 2^n-2i$ for $i \leq 2^{n-1}-4$ and $j = 4, 5, 6$. This shows that hh^σ can be written as

$$hh^\sigma = 1 + (1 + a)^{2^n-(2i+2)} + (1 + a)^{2^n-(2i+1)} + y_\sigma$$

for some $y_\sigma \in A^{2^n-2i}(\mathbb{F}_2C)$. Thus

$$hh^\sigma(1 + a)^{2i} = (1 + a)^{2i} + (1 + a)^{2^n-2} + (1 + a)^{2^n-1}$$

and we have $h \notin H_i^\sigma$. But the equality $hh^\sigma(1 + a)^{2i+2} = (1 + a)^{2i+2}$ states that $h \in H_{i+2}^\sigma$ and this proves that

$$H_0^\sigma \subset H_2^\sigma \subset H_4^\sigma \subset \cdots \subset H_i^\sigma \subset \cdots \subset H_{2^{n-1}-2}^\sigma = V(\mathbb{F}_2C).$$

Now we shall determine the order of the group H_0^σ . The σ -unitary subgroup $V_\sigma(\mathbb{F}_2C)$ of $V(\mathbb{F}_2C)$ is a subgroup of H_0^σ and we shall use the following subgroup

$$J^\sigma = \{ zz^\sigma \mid z \in V(\mathbb{F}_2C), zz^\sigma \in \mathbb{F}_2C^2 \}.$$

According to σ we distinguish two cases. First let σ be the $*$ -involution. We shall prove that $J^* = S_*(C)^2$ and its order is $2^{2^{n-2}-1}$. Indeed, if $u \in J^*$, then $u = zz^* \in \mathbb{F}_2C^2$ for some $z \in V(\mathbb{F}_2C)$ and by Lemma 3,

$$u = zz^* = 1 + \sum_{j=1}^{2^{n-2}-1} \alpha_j(a^{2j} + a^{-2j}).$$

Now the $*$ -symmetric element $y = 1 + \sum_{j=1}^{2^{n-2}-1} \alpha_j(a^j + a^{-j})$ is such that $y^2 = u$. This shows that J^* is a subgroup of $S_*(C)^2$. Conversely, if $u \in S_*(C)^2$, then $u = u^*$ and there exists $y \in S_*(C)$ with $y^2 = u$. Then $yy^* = y^2 = u$ and $u \in J^*$, so $J^* = S_*(C)^2$. The equality of these groups shows that each $y \in S_*(C)$ satisfies

$$y^2 = yy^* = 1 + \sum_{j=1}^{2^{n-2}-1} \alpha_j(a^{2j} + a^{-2j}) \in S_*(C)^2,$$

hence the order of $S_*(C)^2$ is $2^{2^{n-2}-1}$.

The map $\psi_* : V(\mathbb{F}_2 C) \rightarrow S_*(C)$ induces the epimorphism $\psi_* : H_0^* \rightarrow J^*$, and its kernel is the $*$ -unitary subgroup $V_*(\mathbb{F}_2 C)$ and according to (2) $V_*(\mathbb{F}_2 C)$ has order $2^{2^{n-1}+1}$. Thus

$$|H_0^*| = |V_*(\mathbb{F}_2 C)| \cdot |J^*| = |V_*(\mathbb{F}_2 C)| \cdot |S_*(C)|^2 = 2^{3 \cdot 2^{n-2}}.$$

Now, let σ be the $\otimes$ -involution. We shall show that

$$J^{\otimes} = \langle a^{2^{n-1}} \rangle \times S_{\otimes}(C)^2 \quad \text{and} \quad |S_{\otimes}(C)^2| = 2^{\frac{|C|}{4}-1}.$$

For each $y \in S_{\otimes}(C)$ we have

$$y^2 = yy^{\otimes} = 1 + \sum_{j=1}^{2^{n-2}-1} \beta_j (a^{2^j} + a^{-2^j}) \in S_{\otimes}(C)^2,$$

and Lemma 4 asserts that $a^{2^{n-1}} \notin S_{\otimes}(C)^2$ and $|S_{\otimes}(C)^2| = 2^{\frac{|C|}{4}-1} = 2^{2^{n-2}-1}$.

Let $u \in J^{\otimes}$. Then $u = zz^{\otimes} \in \mathbb{F}_2 C^2$ for some $z \in V(\mathbb{F}_2 C)$ and again by Lemma 4 we obtain that

$$\begin{aligned} u = zz^{\otimes} = & \delta_1 (1 + a^{2^{n-1}}) + \\ & + a^{2^{n-1}} \left(1 + \sum_{i=1}^{2^{n-2}-1} \delta_i (a^{2^i+2^{n-1}} + a^{-(2^i+2^{n-1})}) \right), \quad \delta_i \in \mathbb{F}_2. \end{aligned}$$

If $\delta_1 = 0$, then we consider the $\otimes$ -symmetric element

$$\begin{aligned} y_1 = & 1 + \sum_{j \in \{1, 3, \dots, 2^{n-2}-1\}} \delta_j (a^{j+2^{n-2}} + a^{-(j+2^{n-2}+2^{n-1})}) \\ & + \sum_{j \in \{2, 4, \dots, 2^{n-2}-2\}} \delta_j (a^{j+2^{n-2}} + a^{-(j+2^{n-2})}) \in S_{\otimes}(C), \end{aligned}$$

which has the property

$$y_1^2 = 1 + \sum_{j=1}^{2^{n-2}-1} \delta_j (a^{2^j+2^{n-1}} + a^{-(2^j+2^{n-1})}) \in S_{\otimes}(C)^2.$$

Therefore $u = a^{2^{n-1}} y_1^2 \in \langle a^{2^{n-1}} \rangle \times S_{\otimes}(C)^2$.

Now assume that $\delta_1 = 1$. Then the $\otimes$ -symmetric element

$$y_2 = 1 + \sum_{j \in \{1, 3, \dots, 2^{n-1}-1\}} \delta_j (a^j + a^{-j+2^{n-1}}) \\ + \sum_{j \in \{2, 4, \dots, 2^{n-1}-2\}} \delta_j (a^j + a^{-j})$$

is such that $u = y_2^2$ and $u \in J^\otimes \subseteq \langle a^{2^{n-1}} \rangle \times S_\otimes(C)^2$. Conversely, if $u \in \langle a^{2^{n-1}} \rangle \times S_\otimes(C)^2$, then $u = (a^{2^{n-1}})^t d$ for some $d \in S_\otimes(C)^2$. Choose $w \in S_\otimes(C)$ with $w^2 = d$. Then

$$u = (a^{2^{n-1}})^t d = a^t a^{t(2^{n-1}-1)} w^2 = \\ = a^t w a^{t(2^{n-1}-1)} w = a^t w \cdot (a^t w)^\otimes \in J^\otimes.$$

Consequently, $J^\otimes = \langle a^{2^{n-1}} \rangle \times S_\otimes(C)^2$.

Again $\psi_\otimes : V(\mathbb{F}_2 C) \rightarrow S_\otimes(C)$ induces the epimorphism $\psi_\otimes : H_0^\otimes \rightarrow J^\otimes$ and its kernel is the $\otimes$ -unitary subgroup $V_\otimes(\mathbb{F}_2 C)$. This shows that

$$|H_0^\otimes| = |V_\otimes(\mathbb{F}_2 C)| \cdot |J^\otimes| = |V_\otimes(\mathbb{F}_2 C)| \cdot |\langle a^{2^{n-1}} \rangle \times S_\otimes(C)^2| = 2^{3 \cdot 2^{n-2}}.$$

We have established

$$H_0^\sigma \subset H_2^\sigma \subset H_4^\sigma \subset \dots \subset H_{2^{n-1}-2}^\sigma = V(\mathbb{F}_2 C),$$

and $|H_0^\sigma| = 2^{3 \cdot 2^{n-2}}$ from which follows that $[V(\mathbb{F}_2 C) : H_0^\sigma] = 2^{2^{n-2}-1}$ and by the second part of this lemma, the number of different subgroups H_{2l}^σ is $2^{n-2} - 1$. This is possible if and only if $[H_{2l+2}^\sigma : H_{2l}^\sigma] = 2$ and we get that the order of H_{2l}^σ is $2^{3 \cdot 2^{n-2}+l}$ for every $2l < 2^{n-1}$. □

For $0 \leq i < 2^n$ the set

$$L_i^\sigma = \{ h \in V(\mathbb{F}_2 C)[2] \mid (h + h^\sigma)(1 + a)^i = 0 \}$$

is a subgroup of

$$V(\mathbb{F}_2 C)[2] = \left\{ \left(\sum_{j=0}^{2^{n-1}-1} \alpha_j a^j \right) (1 + a)^{2^{n-1}} \in V(\mathbb{F}_2 C) \mid \alpha_j \in \mathbb{F}_2 \right\}. \quad (12)$$

Indeed, each $h_k \in L_i^\sigma$ such that $h_k(1 + a)^i = h_k^\sigma(1 + a)^i$, so

$$h_1 h_2 (1 + a)^i = h_1 h_2^\sigma (1 + a)^i = h_1^\sigma h_2^\sigma (1 + a)^i = (h_1 h_2)^\sigma (1 + a)^i,$$

and L_i^σ is a subgroup. Moreover, easy calculations show that $h^\otimes = h^*$ for each $h \in V(\mathbb{F}_2 C)[2]$. Therefore,

$$L_i^\otimes = L_i^*. \quad (13)$$

Consequently it is sufficient to investigate the properties of L_i^* .

Lemma 10. *The subgroup L_i^* has the following properties:*

1. $L_i^* = V(\mathbb{F}_2 C)[2]$ for $i \geq 2^{n-1}$.
2. For even indices $i = 2l < 2^{n-1}$ the subgroups L_{2l}^* satisfy

$$L_0^* \subset L_2^* \subset L_4^* \subset \cdots \subset L_{2l}^* \subset \cdots \subset L_{2^{n-1}-2}^* = V(\mathbb{F}_2 C)[2],$$

and the order of L_{2l}^* is $2^{2^{n-2}+1+l}$.

3. For odd index $i = 2l + 1 < 2^{n-1}$ the subgroup L_{2l}^* coincides with L_{2l+1}^* .

Proof. First let $i \geq 2^{n-1} - 2$. By (12) each $h \in V(\mathbb{F}_2 C)[2]$ can be represented in the form

$$h = 1 + \alpha_{2^{n-1}}(1+a)^{2^{n-1}} + \alpha_{2^{n-1}+1}(1+a)^{2^{n-1}+1} + u$$

for some $u \in A^{2^{n-1}+2}(\mathbb{F}_2 C)$. The formula (10) asserts

$$h^* = 1 + \alpha_{2^{n-1}}(1+a)^{2^{n-1}} + \alpha_{2^{n-1}+1}(1+a)^{2^{n-1}+1} + v$$

for some $v \in A^{2^{n-1}+2}(\mathbb{F}_2 C)$ and $h + h^* = u + v \in A^{2^{n-1}+2}(\mathbb{F}_2 C)$. But $A^{2^{n-1}+2}(\mathbb{F}_2 C) \subseteq \text{Ann}((1+a)^i)$ for all $i \geq 2^{n-1} - 2$, and we conclude that $(h + h^*)(1+a)^i = 0$. Therefore $L_i^* = V(\mathbb{F}_2 C)[2]$ for all $i \geq 2^{n-1}$ or $i = 2^{n-1} - 2$.

Each symmetric unit $h \in L_0^*$ can be written as

$$h = 1 + \left(\alpha_0 + \alpha_{2^{n-2}} a^{2^{n-2}} + \sum_{j=1}^{2^{n-2}-1} \alpha_j (a^j + a^{-j+2^{n-1}}) \right) (1+a)^{2^{n-1}},$$

where $\alpha_j \in \mathbb{F}_2$. This shows that h has $2^{n-2} + 1$ independent coefficients, thus the order of L_0^* is $2^{2^{n-2}+1}$.

Now we verify that L_{2l}^* is a proper subgroup of $L_{2(l+1)}^*$ for all $2l < 2^{n-1} - 2$. According to (10) for $u = 1 + (1+a)^{2^n-(2l+3)}$ we have

$$u^* = 1 + (1+a)^{2^n-(2l+3)} + (1+a)^{2^n-(2l+2)} + z,$$

where $z \in A^{2^n-(2l+1)}(\mathbb{F}_2 C)$. This shows that $u + u^* = (1+a)^{2^n-(2l+2)} + z$, so

$$(u + u^*)(1+a)^{2l} = (1+a)^{2^n-2} + (1+a)^{2l} z \neq 0,$$

and $u \notin L_{2l}$. But $(u + u^*)(1+a)^{2l+2} = (1+a)^{2^n} = 0$, this means that $u \in L_{2l+2}^*$ and L_{2l}^* is a proper subgroup of L_{2l+2}^* .

Clearly $|V(\mathbb{F}_2 C)[2]| = 2^{2^{n-1}}$, and $[V(\mathbb{F}_2 C)[2] : L_0^*] = 2^{2^{n-2}-1}$. As we saw above the number of different subgroups L_{2l}^* is $2^{n-2} - 1$. The only possibility is that $[L_{2l+2}^* : L_{2l}^*] = 2$ and L_{2l}^* has order $2^{2^{n-2}+1+l}$ for all $2l < 2^{n-1}$.

Finally, we use again the element $u = 1 + (1+a)^{2^n-(2l+3)}$. We note that

$$(u + u^*)(1+a)^{2l+1} = (1+a)^{2^n-1} \neq 0,$$

so $u \notin L_{2l+1}^*$ but $u \in L_{2l+2}^*$. It is easy to see that $L_{2l}^* \subseteq L_{2l+1}^* \subset L_{2l+2}^*$ and $[L_{2l+2}^* : L_{2l}^*] = 2$, so L_{2l}^* coincides with L_{2l+1}^* . □

Proof of the main theorem.

We divide the proof into 3 parts. We begin with the dihedral group. Let us determine the number of the units $x_1 + x_2 b_2$ of type 1 in $V(\mathbb{F}_2 D_{2^{n+1}})$, where $x_2 = 0$ or $x_2 = \gamma(1 + a)^i$, γ is a unit and $i > 0$.

If $x_2 = 0$, then by (5) we have $x_1^2 = 1$. Therefore the number of units of type 1 with $x_2 = 0$ coincides with the order of $V(\mathbb{F}_2 C)[2]$.

Now let $0 < i < 2^{n-1}$ and $x_2 = \gamma(1 + a)^i$. Then

$$x_1^2 = \gamma\gamma^*(a + a^{-1})^i + 1; \quad (14)$$

$$(x_1 + x_1^*)(1 + a)^i = 0. \quad (15)$$

By (14) the element $\gamma\gamma^*(a + a^{-1})^i$ belongs to $\mathbb{F}_2 C^2$, further according to Lemma 8 the number $i = 2l$ is even and

$$H_{2l}^* = \{ \gamma \in V(\mathbb{F}_2 C) \mid \gamma\gamma^*(a + a^{-1})^{2l} \in \mathbb{F}_2 C^2 \}$$

is a subgroup of $V(\mathbb{F}_2 C)$.

For fixed γ and i we determine the number of units $x_1 + x_2 b_2$ of type 1, which satisfy the conditions (14) and (15). If the unit $x'_1 + x_2 b$ is also type 1, then $x'_1 x_1^{-1}$ is a unit of order two and

$$(x'_1 x_1^{-1} + (x'_1 x_1^{-1})^*)(1 + a)^{2l} = x'_1 (x_1^{-1} + (x_1^{-1})^*)(1 + a)^{2l} = 0.$$

Therefore

$$x'_1 x_1^{-1} \in L_{2l}^* = \{ h \in V(\mathbb{F}_2 C)[2] \mid (h + h^*)(1 + a)^{2l} = 0 \},$$

so $x'_1 \in x_1 \cdot L_{2l}^*$ and the number of different elements x'_1 is $|L_{2l}^*|$.

Finally, we shall determine the number of elements $x_2 = \gamma(1 + a)^i$ for a fixed i , where $\gamma \in V(\mathbb{F}_2 C)$. This number coincides with the cardinality of the set

$$K_{2l} = \{ \gamma(1 + a)^{2l} \mid \gamma \in V(\mathbb{F}_2 C), \gamma\gamma^*(a + a^{-1})^{2l} \in \mathbb{F}_2 C^2 \}.$$

Clearly K_{2l} coincides with $H_i^*(1 + a)^{2l}$. But $\gamma(1 + a)^{2l} = \gamma'(1 + a)^{2l}$ if and only if $\gamma^{-1}\gamma'(1 + a)^{2l} = (1 + a)^{2l}$, so $\gamma^{-1}\gamma' \in S_{2l}$. We have established the equality $|K_{2l}| = \frac{|H_{2l}^*|}{|S_{2l}|}$. Hence the number of units of type 1 with the form $x_1 + x_2 b_2$ for a fixed $0 < 2l < 2^{n-1}$ is equal to

$$\frac{|H_{2l}^*|}{|S_{2l}|} \cdot |L_{2l}^*| = 2^{2^n+1}.$$

Now consider the case, when $2^{n-1} \leq i < 2^n$. Then $(a + a^{-1})^i = 0$, and (14) implies that the unit $x_1 + x_2 b_2$ is such that $x_1 \in V(\mathbb{F}_2 C)[2]$ and (15) is always satisfied. Thus $x_2 \in V(\mathbb{F}_2 C)(1 + a)^i$ and there are

$$\frac{|V(\mathbb{F}_2 C)|}{|S_i|} \cdot |V(\mathbb{F}_2 C)[2]|$$

different units of type 1 in $V(\mathbb{F}_2 D_{2^{n+1}})$. We obtain that in $V(\mathbb{F}_2 D_{2^{n+1}})$ the number of units of type 1 is equal to

$$\begin{aligned} & |V(\mathbb{F}_2 C)[2]| + \sum_{l=1}^{2^{n-2}-1} \frac{|H_{2l}^*|}{|S_{2l}|} \cdot |L_{2l}^*| + \sum_{j=2^{n-1}}^{2^n-1} \frac{|V(\mathbb{F}_2 C)|}{|S_j|} \cdot |V(\mathbb{F}_2 C)[2]| = \\ & 2^{2^{n-1}} + \sum_{j=1}^{2^{n-2}-1} 2^{2^{n+1}} + 2^{2^n-1} \sum_{j=2^{n-1}}^{2^n-1} 2^{2^{n-1}} 2^{-j} = 2^{2^n} (2^{n-1} - 1). \end{aligned}$$

Now let us consider the number of units of type 2 in $V(\mathbb{F}_2 D_{2^{n+1}})$. If $x_1 + x_2 b_2$ is a unit of type 2, then x_2 is a unit and according to (5) we have $x_1 = x_1^*$ and $x_2 x_2^* = (1 + x_1)^2$. Evidently $1 + x_1$ is a $*$ -symmetric unit and $x_2 x_2^* \in V(\mathbb{F}_2 C^2)$. For a fixed unit x_2 , by the first part of Lemma 2 the set

$$\{ 1 + x_1 \in S_*(C) \mid (1 + x_1)^2 = x_2 x_2^* \}$$

is a coset of $S_*(C)$ by $S_*(C)[2]$. Therefore the number of the different x_1 is $|S_*(C)[2]|$. Clearly L_0^* coincides with $S_*(C)[2]$, so for a fixed x_2 the number of the different x_1 is $|L_0^*|$.

Since $H_0^* = \{ h \in V(\mathbb{F}_2 C) \mid hh^* \in V(\mathbb{F}_2 C^2) \}$, the number of the different x_2 coincides with $|H_0^*|$. This shows that in $V(\mathbb{F}_2 D_{2^{n+1}})$ the number of the units of type 2 is

$$|H_0^*| \cdot |L_0^*| = 2^{3 \cdot 2^{n-2} + 2^{n-2} + 1} = 2^{2^n + 1}$$

and

$$\Theta_{D_{2^{n+1}}}(2) = 2^{2^n + n - 1} + 2^{2^n}.$$

Now let us consider the generalized quaternion group. Let $x_1 + x_2 b_1$ be a unit of type 2. Similarly to the previous case, x_1 is $*$ -symmetric and

$$x_1^2 + x_2 x_2^* a^{2^{n-1}} = x_1 x_1^* + x_2 x_2^* a^{2^{n-1}} = 1.$$

Since $1 \notin \text{supp}(x_1 x_1^*)$ and $1 \notin \text{supp}(x_2 x_2^* a^{2^{n-1}})$ by Lemma 3 this equality is impossible. Therefore, there is no unit of type 2 in $V(\mathbb{F}_2 Q_{2^{n+1}})$.

By Lemma 7 in $V(\mathbb{F}_2 Q_{2^{n+1}})$ the number of units of type 1 is $(2^{n-1} - 1)2^{2^n}$ and

$$\Theta_{Q_{2^{n+1}}}(2) = 2^{2^n + n - 1} - 2^{2^n}.$$

Finally, let us consider the semidihedral group. According to (13) and Lemma 8, we have $|H_i^{\otimes}| = |H_i^*|$ and $|L_i^{\otimes}| = |L_i^*|$ for each $i < 2^n$. Thus in $V(\mathbb{F}_2SD_{2^{n+1}})$ the number of units of type 1 is $(2^{n-1} - 1)2^{2^n}$ as in the group of units $V(\mathbb{F}_2D_{2^{n+1}})$ for the dihedral group.

Now we shall determine the number of units $x_1 + x_2b_3 \in V(\mathbb{F}_2SD_{2^{n+1}})$ of type 2. Then x_2 is a unit and from (5) it follows that $x_2^{\otimes}x_2 = (x_1 + 1)^2 \in V(\mathbb{F}_2C^2)$ and $(x_1 + x_1^{\otimes})x_2 = 0$. Since x_2 is a unit we get that x_1 is a $\otimes$ -symmetric, so $1 + x_1$ is a $\otimes$ -symmetric unit as well. Evidently $x_2^{\otimes}x_2 = (1 + x_1)^2 \in S_{\otimes}(C)^2$. We have seen that

$$J_{\otimes} = \{ x_2x_2^{\otimes} \mid x_2 \in V(\mathbb{F}_2C), x_2x_2^{\otimes} \in \mathbb{F}_2C^2 \} = \langle a^{2^{n-1}} \rangle \times S_{\otimes}(C)^2,$$

thus the number of different x_2 is

$$\begin{aligned} & \left| \{ x_2 \in V(\mathbb{F}_2C) \mid x_2x_2^{\otimes} \in S_{\otimes}(C)^2 \} \right| = \\ & = |V_{\otimes}(\mathbb{F}_2C)| \cdot |S_{\otimes}(C)^2| = |V_{\otimes}(\mathbb{F}_2C)| \cdot \frac{|J_{\otimes}|}{2} = \frac{|H_0^{\otimes}|}{2}. \end{aligned}$$

For a fixed unit x_2 the set $\{ 1 + x_1 \in S_{\otimes}(C) \mid (1 + x_1)^2 = x_2x_2^{\otimes} \}$ is a coset of $S_{\otimes}(C)$ by $S_{\otimes}(C)[2]$ as the first part of Lemma 2 asserts. Clearly $L_0^{\otimes}$ coincides with $S_{\otimes}(C)[2]$ so for a fixed x_2 the number of the different x_1 is $|L_0^{\otimes}|$. We obtain that the number of units of type 2 is

$$\frac{|H_0^{\otimes}|}{2} \cdot |L_0^{\otimes}| = 2^{3 \cdot 2^{n-2} + 2^{n-2}} = 2^{2^n}$$

and $\Theta_{SD_{2^{n+1}}}(2) = 2^{2^n + n - 1}$, the proof is done. □

REFERENCES

- Baginski, C., *Modular group algebras of 2-groups of maximal class*, Communications in algebra **20(5)** (1992), 1229–1241.
- Berman, S. D., *Group algebras of countable abelian p-group*, Publ. Math. (Debrecen) **14** (1967), 365–405.
- Bovdi, A. A.; Sakach, A., I., *Unitary subgroup of the group of units of a modular group algebra of a finite alelian p-group*, Math. Zametki **45(6)** (1989), 23–29.
- Bovdi, A. A.; Sakach, A., II., *A basis for the unitary subgroup of the group of units in a finite commutative group algebra*, Publ. Math. (Debrecen) **46(1–2)** (1995), 97–120.
- Hill, E. T., *The annihilator of radical powers in the modular group ring of a p-group*, Proc. Amer. Math. Soc. **25** (1970), 811–815.
- Jennings, S.A., *The structure of the group ring of a p-group over a modular field*, Trans. Amer. Math. Soc. **50** (1941), 175–185.

INSTITUTE OF MATHEMATICS AND INFORMATICS, COLLEGE OF NYÍREGYHÁZA, SÓSTÓI ÚT 31/B,
H-4410 NYÍREGYHÁZA, HUNGARY
BALOGHZS@NYF.HU

INSTITUTE OF MATHEMATICS, UNIVERSITY OF DEBRECEN, H-4010 DEBRECEN, P.O. Box 12,
HUNGARY
BODIBELA@MATH.KLTE.HU